Rx only

## **AMINOSYN™II 10%**

An Amino Acid Injection, Sulfite-Free

## Pharmacy Bulk Package — Not for Direct Infusion.

EACH 100 mL CONTAINS: TOTAL AMINO ACIDS APPROX. 10 g. MAY CONTAIN SODIUM HYDROXIDE FOR pH ADJUSTMENT.

ESSENTIAL AMINO ACIDS/100 mL: ISOLEUCINE 660 mg; LYSINE (AS ACETATE SALT) 1050 mg; METHIONINE 172 mg; PHENYLALANINE 298 mg; THREONINE 400 mg; TRYPTOPHAN 200 mg; VALINE 500 mg. NONESSENTIAL AMINO ACIDS/100 mL: N-ACETYL-L-TYROSINE 270 mg; ALANINE 993 mg; ARGININE 1018 mg; GLYCINE 500 mg; PROLINE 722 mg; HISTIDINE 300 mg; SERINE 530 mg; L-ASPARTIC ACID 700 mg; L-GLUTAMIC ACID 738 mg.

ELECTROLYTES (mEq/L): SODIUM 38; ACETATE 71.8. 840 mOsmol/L (ACTUAL). pH 5.8 (5.0 to 6.5) SP. GR.=1.03 FOR IV USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC. SEE INSERT FOR COMPLETE PRODUCT INFORMATION FOR USE OF THE PHARMACY BULK PACKAGE.

DATE ENTERED:

TIME OF ENTRY:

CAUTION: USE ONLY IN LAMINAR FLOW HOOD. ONCE THE OUTLET SITE HAS BEEN ENTERED, THE WITHDRAWAL OF CONTAINER CONTENTS SHOULD BE PROMPTLY COMPLETED IN ONE CONTINUOUS OPERATION. DISCARD CONTAINER NOT LATER THAN 4 HOURS AFTER INITIAL CLOSURE PUNCTURE. SEE INSERT, ADDITIVES MAY BE INCOMPATIBLE WITH FLUID WITHDRAWN FROM THIS CONTAINER. CONSULT WITH PHARMACIST, IF AVAILABLE. WHEN COMPOUNDING ADMIXTURES, USE ASEPTIC TECHNIQUE, MIX THOROUGHLY, DO NOT STORE SOLUTIONS CONTAINING ADDITIVES. BECAUSE OF THE POTENTIAL FOR LIFE-THREATENING EVENTS, CAUTION SHOULD BE TAKEN TO ENSURE THAT PRECIPITATES HAVE NOT FORMED IN ANY PARENTERAL NUTRIENT MIXTURE. STORE AT 20 to 25°C (68 to 77°F). [SEE USP CONTROLLED ROOM TEMPERATURE.] PROTECT FROM FREEZING. AVOID EXPOSURE TO LIGHT. USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. NOT BE USED **SERIES** MUST IN CONNECTIONS. THIS PRODUCT CONTAINS N0 MORE 25 mcg/L OF ALUMINUM.





IM-4371

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA

1*75*0 —

1*5*00 —

1250 **—** 

1000 —

750 **—** 

500 **—** 

250 **—**