

2000 mL

NDC 0990-7171-17

AMINOSYN[®] II

15%

An Amino Acid Injection, Sulfite-Free

Pharmacy Bulk Package—Not for Direct Infusion.

EACH 100 mL CONTAINS: TOTAL AMINO ACIDS APPROX. 15 g. MAY CONTAIN SODIUM HYDROXIDE FOR pH ADJUSTMENT.

ESSENTIAL AMINO ACIDS/100 mL: ISOLEUCINE 990 mg; LEUCINE 1500 mg; LYSINE (AS ACETATE SALT) 1575 mg; METHIONINE 258 mg; PHENYLALANINE 447 mg; THREONINE 600 mg; TRYPTOPHAN 300 mg; VALINE 750 mg.

NONESSENTIAL AMINO ACIDS/100 mL: N-ACETYL-L-TYROSINE 405 mg; ALANINE 1490 mg; ARGININE 1527 mg; GLYCINE 750 mg; PROLINE 1083 mg; HISTIDINE 450 mg; SERINE 795 mg; L-ASPARTIC ACID 1050 mg; L-GLUTAMIC ACID 1107 mg.

ELECTROLYTES (mEq/LITER): SODIUM 50.0, ACETATE 107.6. 1270 mOsmol/L (ACTUAL). pH 5.8 (5.0 to 6.5) SP.GR.=1.05 FOR I.V. USE. USUAL DOSAGE: SEE INSERT. STERILE, NONPYROGENIC. SEE INSERT FOR COMPLETE PRODUCT INFORMATION FOR USE OF THE PHARMACY BULK PACKAGE.

DATE ENTERED: _____

TIME OF ENTRY: _____

AVOID EXPOSURE TO LIGHT.

CAUTION: USE ONLY IN LAMINAR FLOW HOOD. **ONCE THE OUTLET SITE HAS BEEN ENTERED, THE WITHDRAWAL OF CONTAINER CONTENTS SHOULD BE PROMPTLY COMPLETED IN ONE CONTINUOUS OPERATION.** DISCARD CONTAINER NOT LATER THAN 4 HOURS AFTER INITIAL CLOSURE PUNCTURE. SEE INSERT. ADDITIVES MAY BE INCOMPATIBLE WITH FLUID WITHDRAWN FROM THIS CONTAINER. CONSULT WITH PHARMACIST, IF AVAILABLE. WHEN COMPOUNDING ADMIXTURES, USE ASEPTIC TECHNIQUE. MIX THOROUGHLY. DO NOT STORE SOLUTIONS CONTAINING ADDITIVES. BECAUSE OF THE POTENTIAL FOR LIFE-THREATENING EVENTS, CAUTION SHOULD BE TAKEN TO ENSURE THAT PRECIPITATES HAVE NOT FORMED IN ANY PARENTERAL NUTRIENT MIXTURE.

STORE AT 20 TO 25°C (68 TO 77°F). [SEE USP CONTROLLED ROOM TEMPERATURE.] PROTECT FROM FREEZING.

USE ONLY IF SOLUTION IS CLEAR AND CONTAINER IS UNDAMAGED. MUST NOT BE USED IN SERIES CONNECTIONS.

THIS PRODUCT CONTAINS NO MORE THAN 25 mcg/L OF ALUMINUM.

RX ONLY



IM-4370

CONTAINS DEHP

ICU Medical, Inc., Lake Forest, Illinois, 60045, USA

1750 —

1500 —

1250 —

1000 —

750 —

500 —

250 —



icumedical